

Introduction to Riemannian Manifolds

Mica Li


University of California, San Diego

DRP 2024, Mentor: Robert Koirala


Smooth Manifolds

Definition: Smooth Manifold

A **smooth manifold** is a manifold that is endowed with some "smooth structure"




(a) A coordinate chart




(b) A transition map

Examples of Smooth Manifolds



(c) Sphere



(d) Torus

Figure: Examples of smooth manifolds

Geometry on a Manifold?

Question

How do we do geometry on manifolds?

- Length is well-defined on \mathbb{R}^n

Length in \mathbb{R}^n

$$\int_a^b \sqrt{|v(t)|} dt$$

- The issue is that transition maps do not preserve length or angles in general

Idea

We need a new structure so that we can measure length and angles on a manifold

Length on a Manifold?

Question

How do we develop a notion of distance on a manifold?

- In general length is velocity multiplied by time(i.e. vt)
- We need to be able to find the velocity of a curve on a manifold
- Need to define length of a tangent vector

Tangent Spaces

Definition: Tangent Space

The **tangent space** at a point $p \in M$, $T_p M$, is the space of all tangent vectors at that point p .




Figure: Tangent space at a point p

Definition: Riemannian Metrics

A **Riemannian metric** is a "smoothly" chosen inner product on $T_p M$.

An example of the Riemannian metric is

- the standard inner product on \mathbb{R}^n

Length and Distance on a Manifold

Now that we are equipped with the Riemannian metric we can define length and distance

Definition

Let $\gamma : [a, b] \rightarrow M$ be a curve. Then the length of γ is

$$L_g(\gamma) = \int_a^b |\gamma'(t)|_g dt$$

Equipped with this length we can now define the distance metric on a Riemannian manifold.

Distance on a Riemannian Manifold

Let $P = \{\gamma : \gamma \text{ a curve from } p \text{ to } q, p, q \in M\}$. Then,

$$d_g(p, q) = \inf_{\gamma \in P} L_g(\gamma)$$

Definition: Riemannian Manifold

A Riemannian manifold is a pair (M, g) where M is a smooth manifold and g is a Riemannian metric defined on it

An immediate result from this is that

Proposition

Every smooth manifold will admit a Riemannian metric

Where do we go from here?

Using these ideas, one can also generalize

- angles
- volume
- integration and differentiation

Thus, we are now well-equipped to do analysis on manifolds

Acknowledgements

First and foremost I would like to thank my mentor Robert for guiding me through these ten weeks and I am massively appreciative for his patience and guidance. I would also like to thank Prof. Bucur for giving us this opportunity.